

United Nations Children's Pull Ethispis Country Office P.O. Box 1159 Autor Ababa, Ethiopia explane 251-011-0515158 actimite 251-011-0515025 251-011-0517111

HS/120/07

August 15, 2007

Robert Mather Against Malaria Foundation England

Dear Mr. Mather,

URCET would like to discretely bank SWH for its support to provide the propole of Ankenis with long Lating Insocride Metric ULINO to protect themsensive from making. The project has been successful and integrated wave well which the Mattic Enclusion Program USES for [15] and metric programs of the support of the support of the support treatment (Mess Description Program USES) for [15] and metric programs (Mess Description Provide Ankenia a curve planch for the support (nor of the first in this[ba]) and the support programs (Mess Description Provide Ankenia a curve planch for the support (nor of the SWH).

What is of special significance in the Anketha kebeles, is this is one of the first communities to necerbe binark coweraje with all three of Ebbogus main materix costolial activations. Instity, May and an analysis affect communities in Anketaan. Every person (except for pregnant women and infanct) were presumptively treated for malaria using the ArT, Courtem Anternetistre-Limitation (Internetistic). And and and the material Anternetistre-Limitation (Internetistic) and the material Anternetistre-Limitation (Internetistic).



For every child Health: Education, Equality, Protection MOVANCE HUMANITY Secondly all homes were spraved (Indoor Residual Spraving - IRS) to kill all potential malaria vectors in these communities to reduce malaria re-infection after MDA. Thirdly, all families were provided with, on average, two LLINs to protect themselves from re-infection, after having been cleared of parasites during the MDA campaign, IRS will provide high impact vector control, while LLINs are the longer term more cost effective technique for ensuring that the people in these communities remain free from malaria infection for many years more.

We believe that this triple malaria control approach, where treatment (cure) and prevention are tightly integrated. Is likely to substantially reduce malaria in the long term. Of course there is need to assess this by conducting monitoring and evaluation and further studies. To this end the Ethiopian malaria partnership will conduct a Malaria Indicator Survey (MIS) in October 2007, which will provide the most comprehensive data and information ever on the impact of Ethiopia's malaria control program on malaria.

Please finds below table for the number of SWIM malaria ITNs distributed in kebeles of Ankesha Woreda in Amhara region.

List of kebele	Number of ITNs distributed
Chiba	190
Shumata	170
Dugli	180
Ayehu	182
Ambella	100
Kuper	170
Kunzela	1000
Huletu Chaja	980
Chaja Kibirta	1650
Umbiri	820
Bavna	2040
Dimama	1240
Buya	1718
Sostu Segno	160
Total	10,600

Please also find herewith enclosed CD and DVD indicating ITNs distribution and other malaria prevention and control interventions in Ankesha woredas

Best Regar

R.J. NA

Rory Nefrth Project Officer - Health